

# Effect of Different Concentrations of Leukemia Inhibitory Factor on Gene Expression of Vascular Endothelial Growth Factor-A in Trophoblast Tumor Cell Line

Khodakaram Jahanbin, M.Sc.<sup>1</sup>, Mehri Ghafourian, Ph.D.<sup>1,2\*</sup>, Mohammad Rashno, Ph.D.<sup>1,3</sup>

1. Department of Immunology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran  
2. Fertility, Infertility, and Perinatology Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran  
3. Cellular and Molecular Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

## Abstract

**Background:** Several studies have shown that leukemia inhibitory factor (LIF) is one of the most important cytokines participating in the process of embryo implantation and pregnancy, while, the role of this factor on vascular endothelial factor-A (VEGF-A), as one of the most important angiogenic factor, has not been fully investigated yet. The aim of this study was to evaluate the effect of LIF on gene expression of *VEGF* in the choriocarcinoma cells (JEG-3).

**Materials and Methods:** In this experimental study, JEG-3 choriocarcinoma cells were treated with different concentrations of LIF (1, 10, and 50 ng) for 6, 12, 24, 48 and 72 hours. Expression of *VEGF* was analyzed by real-time PCR. Delta CTs were subjected to one-way analysis of variance (ANOVA) and a post hoc Tukey's test by SPSS version 25.0 software for data analyzing.

**Results:** In the stimulated cells, different concentrations of LIF caused significant decrease of *VEGF* gene expression ( $P<0.05$ ) at 12, 24 and 48 hours. In contrast, it was increased after 72 hours ( $P<0.001$ ). Analysis of data after 6 hours also showed that level of *VEGF* gene expression was significantly decreased by increasing LIF concentration ( $P<0.001$ ).

**Conclusion:** Expression level of *VEGF* gene was decreased in trophoblast cells (except after 72 hours) under the effect of different concentrations of LIF in a time-dependent manner. So, this study showed that further studies are needed to determine the effect of LIF on other angiogenic factors in trophoblast cells.

**Keywords:** Leukemia Inhibitory Factor, Trophoblast, Vascular Endothelial Growth Factor-A

**Citation:** Jahanbin Kh, Ghafourian M, Rashno M. Effect of different concentrations of leukemia inhibitory factor on gene expression of vascular endothelial growth factor-A in trophoblast tumor cell line. *Int J Fertil Steril.* 2020; 14(2): 116-121. doi: 10.22074/ijfs.2020.6058.

This open-access article has been published under the terms of the Creative Commons Attribution Non-Commercial 3.0 (CC BY-NC 3.0).

## Introduction

Leukemia inhibitory factor (LIF) is a glycoprotein cytokine with a molecular weight of 38-67 kDa. That is a member of the interleukin 6 family. LIF receptor is a heterodimer composed of two chains, gp130 and leukemia inhibitory factor receptor- $\beta$  (LIFR- $\beta$ ) expressing on the surface of trophoblast cells (1, 2). LIF induce tyrosine phosphorylation in signal transducers and transcription factors of several trophoblast cell types, like choriocarcinoma cell line (JEG-3) (3, 4). Phosphorylation and signal transduction lead to migration, invasion, stimulation or suppression of various categories of genes in trophoblast cells (5, 6). Janus kinase 1 (JAK-1) and Signal transducer and activator of transcription 3 (STAT-3), play important roles in the signal transduction factors and activation of transcription in the LIF signaling (7, 8). VEGF is a homodimer glycoprotein which can stimulate angiogenesis and vasculogenesis by two types of its receptors like Fms-

like tyrosine kinase 1 (Flt1) and kinase insert domain receptor (KDR) (9, 10). VEGF has many roles in early pregnancy, such as oocyte maturation and development, trophoblast proliferation, placenta angiogenesis, embryo implantation, maternal blood vessel growth and development of the embryonic blood vessels (11). Formation of the placenta in uterus depends on differentiation of extravillous cytotrophoblast (EVT) for invasion to the uterine stroma and forming endovascular trophoblast (12, 13). Incorrect differentiation of EVT cells leads to disruption of spiral artery remodeling, and this impairment in spiral artery remodeling can lead to preeclampsia and defective development of the fetus (13). Trophoblast invasion is a localized and temporary process. That is the main factor in the regulation of implantation and supply of oxygen to the fetus. By *VEGF* gene inactivation, invasion and migration of trophoblast cells are reduced (14). VEGF-A is one of the main factors of EVT differentiation to the

Received: 1/August/2019, Accepted: 11/December/2019

\* Corresponding Address: P.O.Box: 6135715794, Department of Immunology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran  
Email: ghafourianbm@gmail.com



Royan Institute  
International Journal of Fertility and Sterility  
Vol 14, No 2, July-September 2020, Pages: 116-121

endovascular trophoblast (15). Anti-angiogenic factors that reduce the amount of VEGF-A is one of the factors inhibiting formation of spiral arteries, which eventually associated with the creation of preeclampsia (16). VEGF-A is one of the factors encoded by *VEGF* gene. Studies have shown that among all growth factors encoded by this gene, VEGF-A is the most potent type in stimulating angiogenesis (17). During formation of placenta, EVT, involving in vascular reconstruction, acquire the features associated with epithelial cells, following the production of VEGF and its receptor expression on the surface (12, 18). These cells migrate to decidua, followed by replacement of the endothelial cells in the spiral arteries to form spiral arteries (19). In this study, a choriocarcinoma cell line JEG-3 (derived from fetal trophoblast tumor) was used as EVTs (20). This cell line has many biological and biochemical features of EVTs (Cells lining the blood vessels of villus in the placenta) (21). This cell line is able to produce progesterone, hCG, steroids and other hormones in the placenta (22). In this study we aimed to evaluate *VEGF* gene expression levels in trophoblast tumor cell line (JEG-3) at different times, while these cells were treated by different concentrations of LIF.

## Materials and Methods

This experimental study approved by Ethics Committee of Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran (Ethics code: IR.AJUMS.REC.1395.577).

### Cell culture and treatment

JEG-3 choriocarcinoma cells were purchased from the Pasteur Institute of Iran (Tehran, Iran). These cells were maintained in Dulbecco's modified Eagle's medium-F12 (DMEM-F12; GIBCO, Ireland) supplemented with 10% heat-inactivated fetal bovine serum (FBS; GIBCO, Ireland) along with penicillin (BioIdea, Iran; 100 units/ml) and streptomycin (BioIdea; 100 µg/ml). All JEG-3 cultures were commenced at 106 cells/175-cm<sup>2</sup> flask and maintained under standardized conditions (37°C, 5% CO<sub>2</sub>, humidified atmosphere). The cells were trypsinized twice a week when confluence was estimated at over 75%. For all assays, JEG-3 cells were adjusted to 10<sup>5</sup> cells/ml. The cells (10<sup>5</sup> cell/ml) were seeded in six-well plates, following the resuspension in complete growth media. Before adding the stimuli, the cells were starved for 2 hours in medium without FBS. The cells were cultured per well in the presence and absence of different concentrations (1 ng/ml, 10 ng/ml, 50 ng/ml) (23, 24) of human LIF (Sigma-Aldrich, Germany), while non-stimulated cells were

included as controls. Treated and non-treated cells were incubated for 3, 6, 12, 24, 48 and 72 hours at 37°C with 5% CO<sub>2</sub>. The cell culture supernatants were then collected by aspiration and centrifugation at 1000 g for 5 minutes and they were stored at -70°C until cytokine analysis. JEG-3 cells were harvested and kept at -70°C until total ribonucleic acid (RNA) extraction.

### Ribonucleic acid (RNA) isolation and real-time polymerase chain reaction (PCR) analysis

RNA was isolated using TRI Reagent (SinaClonCo., Iran). According to the manufacturer's protocol, and the purity of extracted RNA was determined by the A260/A280 ratio (A260/A280 ratio was 1.8). 50-100 ng RNA was reverse transcribed using cDNA synthesis kit (SinaClonCo.) and relative changes in *VEGF* mRNA level was quantified by real-time reverse transcription PCR (RT-PCR). Expression level of *VEGF* was determined by quantitative RT-PCR (qRT-PCR) using SYBR Green® Premix Ex Taq (Takara, Japan) dye detection method on ABI StepOne PCR instrument (Applied Biosystems, USA), compared to *GAPDH* as an internal control. Initial denaturation at 95°C for 10 minutes, 40 cycles of annealing at 95°C for 15 seconds and extension at 68°C for 60 seconds. Rest 2009 and Excel software were used for the analysis of gene expression ratio. Gene-specific primers for *VEGF* and *GAPDH* are summarized in Table 1. The fold change for target genes normalized by internal control was determined by the formula  $2^{-\Delta\Delta Ct}$ . All reactions were run in duplicate.

### Statistical analysis


All of the experiments were repeated in triplicates and data were demonstrated as means ± standard error (SE). Statistical software SPSS 25.0 and Graphpad Prism 8.0.1 were used for data analysis. Delta CTs were subjected to one-way ANOVA and a post hoc Tukey's test, while the non-parametric Kruskal-Wallis test was used to compare the results of different experimental days. P values lower than 0.05 were considered statistically significant.

## Results


Effects of different concentrations of LIF on VEGF gene expression level This study evaluated the effects of different concentrations of LIF on *VEGF* gene expression in different time periods, compared to untreated cells. The results are described (Fig.1, 2) in more details.

**Table 1:** Gene specific primers used for RT-PCR

Primer (accession)	Sequence (5'-3')	T <sub>m</sub>	Amplicon size
<i>VEGF</i> (NM_001287044.1)	F: AGGAGGAGGGCAGAATCATCA R: CTCGATTGGATGGCAGTAGCT	60	76 bp
<i>GAPDH</i> (NM_002046.5)	F: TGGGCTACACTGAGCACCAG R: CAGCGTCAAAGGTGGAGGAG	60	72 bp



**Fig.1:** VEGF gene expression level at different time points, under treatment with different concentrations of LIF. The effect of different concentrations of LIF (1, 10 and 50 ng) on VEGF gene expression after A. 6 hours; B. 12 hours; C. 24 hours; D. 48 hours; and E. 72 hours. Cells that did not treated by LIF were considered at any time as control, and the VEGF gene expression was measured in treated cells relative to these untreated cells. \*, P<0.05, \*\*, P<0.01, and \*\*\*, P<0.001



**Fig.2:** Comparing total *VEGF* gene expression at different time (6, 12, 24, 48 and 72 hours) under the treatment of different concentrations of LIF. \*;  $P < 0.05$ , \*\*;  $P \leq 0.01$ , \*\*\*;  $P \leq 0.001$ , and h; Hours.

### Six hours treatment

An analysis of 6 hours data showed that by increasing LIF concentration, level of *VEGF* gene expression was decreased. In this time point, there is a significant difference ( $P < 0.001$ ) between the rate of *VEGF* gene expression in comparison with each other at different concentrations and control (Fig.1A).

### Twelve hours treatment

After 12 hours, there was a significant reduction in the *VEGF* gene expression in all three concentrations of LIF treatment than control ( $P < 0.001$ ). The lowest *VEGF* gene expression level was observed at 1 ng concentration of LIF. The results of 10 ng concentration of LIF were almost similar to the 1 ng (the difference between 1 and 10 ng was not significant). At 50 ng concentration of LIF, *VEGF* expression level was higher than the both concentrations of 1 and 10 ng ( $P < 0.05$ , Fig.1B).

### Twenty-four hours treatment

Twenty-four hours after cells treatment with different concentrations of LIF, the results showed lowest expression of the *VEGF* gene at the concentration of 10 ng ( $P < 0.001$ ). Using 10 ng ( $P < 0.001$ ) and 50 ng ( $P < 0.01$ ) concentrations, there was a significant decrease in gene expression compared to control, but at 1 ng concentration, there was no significant decrease in the gene expression ( $P = 0.324$ ). Comparing gene expression between different concentrations of LIF showed a significant difference between the concentration of 1 ng and 10 ng ( $P = 0.004$ , Fig.1C).

### Forty-eight hours treatment


After 48 hours, like 12 and 24 hours, *VEGF* gene expression was decreased by treating with different concentrations of LIF, compared to control ( $P < 0.001$ ), and the lowest gene expression was observed at 1 ng in comparison with 10 ng ( $P < 0.01$ ) and 50 ng ( $P < 0.05$ ). *VEGF* gene expression was more in 10 ng than the other two concentrations (1 and 50 ng) of LIF ( $P < 0.01$ ; Fig. 1D).

### Seventy-two hours treatment

After 72 hours, effect of LIF on the *VEGF* gene expression was reversed, and contrary to the previous times, in all three concentrations of LIF, we observed a dramatic increased expression of the *VEGF* gene, in comparison with control ( $P < 0.001$ ; Fig.2). The maximum *VEGF* gene expression was observed at 10 ng of LIF concentration, which had significant difference in comparison with 1 ng concentration of LIF ( $P < 0.05$ ). But, the difference between 50 ng and 10 ng LIF concentrations was not significant (Fig.1E).

### VEGF gene expression at different time points

As shown in Figure 3, *VEGF* gene expression was dramatically decreased ( $P < 0.001$ ) at 12, 24, and 48 hours after cell treatment with LIF, in comparison with 6 hours treatment. In contrast to decrease in the *VEGF* gene expression at 12, 24 and 48 hours, we determined a significant increase ( $P < 0.001$ ) in *VEGF* gene expression at 72 hours compared to other time points.



**Fig.3:** Changes in *VEGF* gene expression at different time points of inducing LIF to the cells. \*;  $P < 0.05$ , \*\*;  $P \leq 0.01$ , \*\*\*;  $P \leq 0.001$ .

### Discussion

Pregnancy is a complex process that depends on many factors. Studies have shown that cytokines, growth factors and several transcription factors play important roles in embryo implantation. For example, production of LIF by endometrial cells is essential for the beginning of implantation (25, 26). Data obtained from mice and humans have shown that among the all molecules expressed in uterus, LIF plays the most important role in embryo implantation (27). Formation of new blood vessels is called angiogenesis and it accompanies with migration, growth and differentiation of endothelial cells (28). Angiogenesis usually occurs during the menstrual cycle or estrus to convert the ovulation follicles to corpus luteum which leads to the synthesis of progesterone and restructure of the endometrium. This culminates in maintenance of embryo implantation (29). Angiogenesis and vasculogenesis are essential processes for increasing blood flow to the fetus and, consequently, supplying the nutrients and oxygen needed by fetus (13, 30, 31). Several growth factors con-

control angiogenesis and vasculogenesis during pregnancy. Among these factors, VEGF plays a critical role in the development of the placenta and formation of vessels. Carmeliet et al. (32) showed that deactivation of only one VEGF allele leads to fetal death through angiogenesis disruption. Shalaby et al. (33) by disrupting Vascular endothelial growth factor receptor 1 (VEGFR1), Fong et al. (34) by disrupting VEGFR2 and Tsoi et al. (35) by disruption of neuropilin-1 and -2 (all of them are VEGF receptors) determined similar results to Carmeliet et al. (32). Adequate blood supply to the placenta is highly dependent on regulated invasion and trophoblast vascular remodeling in uterus (36, 37). Extravillous trophoblast (EVT) is a subset of trophoblasts that play the most important role in invasion (the same mechanism as cancerous cells for invasion) to the mother's uterus and vascular remodeling. This eventually acquires the phenotype of endothelial cells and improves artery formation (37). Previous studies have shown that EVTs have receptors for VEGF at their surface and message through these receptors which stimulate invasion, switch phenotype to endovascular cells and tube formation in EVT cells (37, 38). Defects in EVT invasion and angiogenesis have been observed in disorders, such as preeclampsia and intrauterine growth restriction (IUGR) (37). Due to the vital role of vascular formation by trophoblast cells (especially EVTs) in pregnancy and implantation, in this study, we decided to investigate the effect of LIF on one of the most important angiogenic factors, VEGF, in EVTs. For this purpose, we had to select an appropriate cell line with similar features to EVT cells. According to the previous studies (20, 39), JEG-3 cell line was selected. The results of this study showed that LIF could have a dual effect on *VEGF* gene expression with respect to time. So that at 12, 24, and 48 hours, *VEGF* gene expression was decreased, while it was increased at 6 and 72 hours (the increase of *VEGF* gene expression at 6 hours depended on the concentration of LIF showing a significant decrease at 50 ng concentration of LIF in contrast to 1 and 10 ng).

Considering the mentioned roles for VEGF during pregnancy and relevant disorders, as well as the important role of LIF during pregnancy, we decided to investigate the effect of LIF on the level of *VEGF* gene expression in trophoblast cells. Regarding the results, it was found that the expression of the *VEGF* gene in trophoblast tumor cells treated by LIF was reduced in concentration- and time-dependent manners. Although expression of the *VEGF* gene was significantly increased after 72 hours, a study has previously shown that half-life of the LIF attachment to its receptor is slightly more than 24 hours (40). It can be concluded that after 72 hours, interactions between LIFs and their receptors are broken-down and the LIF signaling from their receptors are ended in trophoblast cells. As the results of this study showed, different concentrations of LIF can reduce the rate of *VEGF* gene expression depending on the time. So given the fact that *VEGF* gene expression level was decreased in LIF-treated cells, assessment of the production and secretion of

VEGF protein in treated trophoblast cells is vital. Further investigations have to be performed on the other angiogenic factors to clarify the role of LIF on angiogenesis procedure in trophoblast cells.

## Conclusion

In conclusion, recent studies have shown that both LIF and VEGF are essential for maintaining and initiating the pregnancy process. It has also been found that angiogenesis process is a critical procedure in embryonic trophoblast cells for a normal pregnancy. VEGF-A is one of the most important angiogenic factors. Therefore, in this study we investigated the effect of LIF on *VEGF* gene expression in JEG-3 cell line as extravillous trophoblast cells. According to the results of this study, LIF causes a significant decrease in gene expression level of VEGF-A in JEG-3 cells. Further studies are needed to determine the action mechanism of LIF in angiogenesis of trophoblast cells.

## Acknowledgements

Ahvaz Jundishapur University of Medical Sciences has financially supported this study. This article is the result of thesis for Khodakaram Jahanbin to obtain a master degree in Immunology (Grant number: FIRC-9507). There is no conflict of interest in this study.

## Authors' Contributions

M.G., M.R.; Contributed to concept and design and were responsible for overall supervision. K.J.; Contributed to all experimental work, data and statistical analysis, as well as interpretation of data. M.G.; Drafted the manuscript, which was revised by K.J. All authors read and approved the final manuscript.

## References

1. Sun X, Bartos A, Whitsett JA, Dey SK. Uterine deletion of Gp130 or Stat3 shows implantation failure with increased estrogenic responses. *Mol Endocrinol*. 2013; 27(9): 1492-1501.
2. Nicola NA, Babon JJ. Leukemia inhibitory factor (LIF). *Cytokine Growth Factor Rev*. 2015; 26(5): 533-544.
3. Suman P, Malhotra SS, Gupta SK. LIF-STAT signaling and trophoblast biology. *JAKSTAT*. 2013; 2(4): e25155.
4. Morales-Prieto DM, Barth E, Murrieta-Coxca JM, Favaro RR, Gutiérrez-Samudio RN, Chaiwangyen W, et al. Identification of miRNAs and associated pathways regulated by Leukemia Inhibitory Factor in trophoblastic cell lines. *Placenta*. 2019; 88: 20-27.
5. Chen X, Tong C, Li H, Peng W, Li R, Luo X, et al. Dysregulated expression of RPS4Y1 (Ribosomal Protein S4, Y-Linked 1) impairs STAT3 (Signal Transducer and Activator of Transcription 3) signaling to suppress trophoblast cell migration and invasion in preeclampsia. *Hypertension*. 2018; 71(3): 481-490.
6. Massimiani M, Vecchione L, Piccirilli D, Spitalieri P, Amati F, Salvi S, et al. Epidermal growth factor-like domain 7 promotes migration and invasion of human trophoblast cells through activation of MAPK, PI3K and NOTCH signaling pathways. *Mol Hum Reprod*. 2015; 21(5): 435-451.
7. Takahashi Y, Takahashi M, Carpino N, Jou ST, Chao JR, Tanaka S, et al. Leukemia inhibitory factor regulates trophoblast giant cell differentiation via Janus kinase 1-signal transducer and activator of transcription 3-suppressor of cytokine signaling 3 pathway. *Mol*

- Endocrinol. 2008; 22(7): 1673-1681.
8. Murphy MJ, Halow NG, Royer PA, Hennebold JD. Leukemia inhibitory factor is necessary for ovulation in female rhesus macaques. *Endocrinology*. 2016; 157(11): 4378-4387.
  9. Xiao Z, Li S, Yu Y, Li M, Chen J, Wang F, et al. VEGF-A regulates sFit-1 production in trophoblasts through both Fit-1 and KDR receptors. Vascular permeability factor/vascular endothelial growth factor: a critical cytokine in tumor angiogenesis and a potential target for diagnosis and therapy. *Mol Cell Biochem*. 2018; 449(1-2): 1-8.
  10. Dvorak HF. Vascular permeability factor/vascular endothelial growth factor: a critical cytokine in tumor angiogenesis and a potential target for diagnosis and therapy. *J Clin Oncol*. 2002; 4368-43890.
  11. Red-Horse K, Zhou Y, Genbacev O, Prakobphol A, Foulk R, McMaster M, et al. Trophoblast differentiation during embryo implantation and formation of the maternal-fetal interface. *J Clin Invest*. 2004; 114(6): 744-754.
  12. Pollheimer J, Vondra S, Baltayeva J, Beristain AG, Knöfler M. Regulation of placental extravillous trophoblasts by the maternal uterine environment. *Front Immunol*. 2018; 9: 2597.
  13. Maynard SE, Karumanchi SA. Angoing factors and preeclampsia. *Semin Nephrol*. 2011; 31(1): 33-46.
  14. Dubinsky V, Poehlmann TG, Suman P, Gentile T, Markert UR, Gutierrez G. Role of regulatory and angiogenic cytokines in invasion of trophoblastic cells. *Am J Reprod Immunol*. 2010; 63(3): 193-199.
  15. Li Y, Zhu H, Klausen C, Peng B, Leung PC. Vascular endothelial growth factor-A (VEGF-A) mediates activin A-induced human trophoblast endothelial-like tube formation. *Endocrinology*. 2015; 156(11): 4257-4268.
  16. Helmo FR, Lopes AMM, Carneiro ACDM, Campos CG, Silva PB, dos Reis Monteiro MLG, et al. Angiogenic and antiangiogenic factors in preeclampsia. *Pathol Res Pract*. 2018; 214(1): 7-14.
  17. Arutyunyan I, Fatkhudinov T, Kananykhina E, Usman N, Elchaninov A, Makarov A, et al. Role of VEGF-A in angiogenesis promoted by umbilical cord-derived mesenchymal stromal/stem cells: in vitro study. *Stem Cell Res Ther*. 2016; 7: 46.
  18. Babischkin JS, Aberdeen GW, Lindner JR, Bonagura TW, Pepe GJ, Albrecht ED. Vascular endothelial growth factor delivery to placental basal plate promotes uterine artery remodeling in the primate. *Endocrinology*. 2019; 160(6): 1492-1505.
  19. Tessier DR, Yockell-Lelièvre J, Gruslin A. Uterine spiral artery remodeling: the role of uterine natural killer cells and extravillous trophoblasts in normal and high-risk human pregnancies. *Am J Reprod Immunol*. 2015; 74(1): 1-11.
  20. Olivares EG, Muñoz R, Tejerizo G, Montes MJ, Gómez-Molina F, Abadía-Molina AC. Decidual lymphocytes of human spontaneous abortions induce apoptosis but not necrosis in JEG-3 extravillous trophoblast cells. *Biol Reprod*. 2002; 67(4): 1211-1217.
  21. Drwal E, Rak A, Gregoraszczyk EL. Differential effects of ambient PAH mixtures on cellular and steroidogenic properties of placental JEG-3 and BeWo cells. *Reprod Toxicol*. 2019; 86: 14-22.
  22. Hu W, Gao F, Zhang H, Hiromori Y, Arakawa S, Nagase H, et al. Activation of peroxisome proliferator-activated receptor gamma and disruption of progesterone synthesis of 2-ethylhexyl diphenyl phosphate in human placental choriocarcinoma cells: comparison with triphenyl phosphate. *Environ Sci Technol*. 2017; 51(7): 4061-4068.
  23. Fitzgerald JS, Tsareva SA, Poehlmann TG, Berod L, Meissner A, Corvinus FM, et al. Leukemia inhibitory factor triggers activation of signal transducer and activator of transcription 3, proliferation, invasiveness, and altered protease expression in choriocarcinoma cells. *Int J Biochem Cell Biol*. 2005; 37(11): 2284-2296.
  24. Suman P, Gupta SK. STAT 3 and ERK 1/2 cross-talk in leukaemia inhibitory factor mediated trophoblastic JEG-3 cell invasion and expression of Mucin 1 and Fos. *Am J Reprod Immunol*. 2014; 72(1): 65-74.
  25. Cheng J, Rosario G, Cohen TV, Hu J, Stewart CL. Tissue-specific ablation of the LIF receptor in the murine uterine epithelium results in implantation failure. *Endocrinology*. 2017; 158(6): 1916-1928.
  26. Bulun SE. Steroids, cytokines, and implantation. *Endocrinology*. 2017; 158(6): 1575-1576.
  27. Red-Horse K, Zhou Y, Genbacev O, Prakobphol A, Foulk R, McMaster M, et al. Trophoblast differentiation during embryo implantation and formation of the maternal-fetal interface. *J Clin Invest*. 2004; 114(6): 744-754.
  28. Chun CZ, Sood R, Ramchandran R. Vasculogenesis and angiogenesis. In: North PE, Sander T, editors. *Vascular tumors and developmental malformations*. New York: Springer; 2016; 77-99.
  29. Tamanini C, De Ambrogi M. Angiogenesis in developing follicle and corpus luteum. *Reprod Domest Anim*. 2004; 39(4): 206-216.
  30. Lang U, Baker RS, Braems G, Zygmunt M, Künzel W, Clark KE. Uterine blood flow-a determinant of fetal growth. *Eur J Obstet Gynecol Reprod Biol*. 2003; 110(Suppl 1): S55-S61.
  31. Tal R, Dong D, Shaikh S, Mamillapalli R, Taylor HSJBor. Bone-marrow-derived endothelial progenitor cells contribute to vasculogenesis of pregnant mouse uterus. *Biol Reprod*. 2019; 100(5): 1228-1237.
  32. Carmeliet P, Ferreira V, Breier G, Pollefeyt S, Kieckens L, Gertsenstein M, et al. Abnormal blood vessel development and lethality in embryos lacking a single VEGF allele. *Nature*. 1996; 380(6573): 435-439.
  33. Shalaby F, Rossant J, Yamaguchi TP, Gertsenstein M, Wu XF, Breitman ML, et al. Failure of blood-island formation and vasculogenesis in Flk-1-deficient mice. *Nature*. 1995; 376(6535): 62-66.
  34. Fong GH, Rossant J, Gertsenstein M, Breitman ML. Role of the Flt-1 receptor tyrosine kinase in regulating the assembly of vascular endothelium. *Nature*. 1995; 376(6535): 66-70.
  35. Tsoi SC, Wen Y, Chung JY, Chen D, Magness RR, Zheng J. Co-expression of vascular endothelial growth factor and neuropilin-1 in ovine fetoplacental artery endothelial cells. *Mol Cell Endocrinol*. 2002; 196(1-2): 95-106.
  36. Burton GJ, Woods AW, Jauniaux E, Kingdom JC. Rheological and physiological consequences of conversion of the maternal spiral arteries for uteroplacental blood flow during human pregnancy. *Placenta*. 2009; 30(6): 473-482.
  37. Lala PK, Nandi P. Mechanisms of trophoblast migration, endometrial angiogenesis in preeclampsia: The role of decorin. *Cell Adh Migr*. 2016; 10(1-2): 111-125.
  38. Lala N, Girish GV, Cloutier-Bosworth A, Lala PK. Mechanisms in decorin regulation of vascular endothelial growth factor-induced human trophoblast migration and acquisition of endothelial phenotype. *Biol Reprod*. 2012; 87(3): 59.
  39. Zygmunt M, Hahn D, Münstedt K, Bischof P, Lang U. Invasion of cytotrophoblastic JEG-3 cells is stimulated by hCG in vitro. *Placenta*. 1998; 19(8): 587-593.
  40. Hilton DJ, Nicola NA. Kinetic analyses of the binding of leukemia inhibitory factor to receptor on cells and membranes and in detergent solution. *J Biol Chem*. 1992; 267(15): 10238-10247.